



Syddansk Universitet

Glucocorticoid-induced changes in the geometry of osteoclast resorption cavities are sufficient to alter bone quality: an assessment through a model simulating bone stiffness

Vanderoost, J; S  , Kent; Merrild, Ditte Marie Horslev; Delaiss  , Jean-Marie; van Lenthe, G-H

Published in:
  ben Forskerdag

Publication date:
2011

Document Version
Submitted manuscript

[Link to publication](#)

Citation for pulished version (APA):

Vanderoost, J., S  , K., Merrild, D. M. H., Delaiss  , J-M., & van Lenthe, G-H. (2011). Glucocorticoid-induced changes in the geometry of osteoclast resorption cavities are sufficient to alter bone quality: an assessment through a model simulating bone stiffness.   ben Forskerdag, 29.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Download date: 16. jan.. 2017

Glucocorticoid-induced changes in the geometry of osteoclast resorption cavities are sufficient to alter bone quality: an assessment through a model simulating bone stiffness

Forfattere	Vanderoost J ¹ , Sørensen K ² , Merrild DMH ² , Delaissé J-M ² , van Lenthe GH ¹ ¹ Division of Biomechanics and Engineering Design, K.U.Leuven, Leuven, Belgium, ² Department of Clinical Cell Biology, Vejle Hospital, University of Southern Denmark, Vejle, Denmark
E-mail	kent.soree@slb.regionsyddanmark.dk
Foredrag	Kent Sørensen
Formål	<p>Loss of bone strength and prevention of fracture is a major concern for clinicians dealing with bone pathologies, such as postmenopausal osteoporosis and glucocorticoid-induced osteoporosis (GIO). For postmenopausal osteoporosis fracture risk is mostly defined by BMD thereby taking into account deficits in bone mass and degree of mineralization. However, in the case of GIO it is well known that BMD is a poor predictor of fracture risk. We have recently found that glucocorticoid (GC) causes osteoclasts (OCs) to change their resorptive behavior and making large elongated resorption cavities instead of individual circular resorption cavities, but without affecting the total eroded surface.</p> <p>The aim is a change in resorptive behavior of OCs sufficient to increase the fracture risk in GIO.</p>
Metode	The three dimensional characteristics of resorption cavities generated in vitro by human OCs in the presence or absence of GC were recorded. These data were used to model the impact of GC-induced resorption on the mechanical properties of bone using the beam-shell finite element model of trabecular bone.
Resultater	The model showed that resorption cavities generated in the presence of GC caused a statistical significant loss of stiffness as simulated for three bone sites (4 th lumbar spine, femoral neck and iliac crest), when compared to control. This strongly suggests that the altered resorptive behavior of the OC due to GC-exposure is sufficient to enhance the fracture risk at these sites. It is worth noting that the GC-induced changes of the shape of cavitations had the highest impact on bone stiffness in the 4 th lumbar spine. It is therefore interesting to note that GIO is in particular associated to an increased risk of vertebral fractures.
Konklusion	Changes in the shape of OC resorption cavities are sufficient to reduce bone stiffness/strength independent of the extent of eroded surface. This shows the general need of taking the quality of OC bone resorption into account and not only broad quantitative measurements of OC activity as usually done through histomorphometry or bone markers.